
Horseshoe Valley Aquifer

McLean County


Aquifer At-a-Glance				
Area	9.8 square miles			
Aquifer Type	Unconfined Surficial			
Major Land Uses over Aquifer	Grassland/Pasture (59%)			
(percentage of aquifer area covered in 2017) ¹	Crops (34%)			
Depth to Water* 2	7-35 feet			
Total Unique Wells Sampled	8			
Wells Sampled in 2021	3			
Samples Collected in 2021	4			
Years Sampled	1996, 2001, 2006, 2011, 2016, 2021			

*Depths to water may vary seasonally, year to year, and across the aquifer

- Aguifer materials consist of sands and gravels that fill an ancient valley which was carved by meltwater from a glacier.2
- The aquifer ranges from 15-44 feet thick and averages about 30 feet thick.2
- Domestic, irrigation, and stock wells are installed in the aquifer.
- In North Dakota, permits are required to withdraw large quantities of groundwater. In 2020, 108 million gallons of permitted water were drawn from the aguifer; irrigation use consumed the largest quantity of water. For more information on water use and permits, contact the North Dakota Department of Water Resources (dwr.nd.gov).



2020 Horseshoe Valley aquifer permitted water use (from North Dakota Department of Water Resources (dwr.nd.gov))



About the Agricultural Groundwater Monitoring Program

- The North Dakota Department of Environmental Quality monitors a network of wells in approximately 50 surficial aguifers that are at elevated risk of agricultural contamination.
- Aquifers are sampled on a 5-year rotation.
- Monitoring began in 1992.
- The vast majority of these aquifers are located in central and eastern North Dakota.
- Water is tested for 21 general chemistry parameters, eight trace metals, and 64 pesticides.

US Department of Agriculture, 2017, National Agricultural Statistics Service Cropland Data Layer. Klausing, R.L., 1974, Ground-Water Resources of McLean County, North Dakota. North Dakota State Water Commission County Groundwater Studies 19-Part 3; North Dakota Geological Survey Bulletin

Water Chemistry

Is Aquifer	
Water	
High in?	

	Analyte	Result	2021 Median Concentration	Potential Effects
	Arsenic	Locally	0.009 mg/L	Skin or circulatory system damage, increased cancer risk
r	Iron	YES	7.84 mg/L	Metallic taste/odor, discoloration of surfaces
	Manganese	YES	0.69 mg/L	
?	Sodium	NO	30.2 mg/L	Taste, people with certain health conditions may need to limit intake
	Sulfate	NO	150 mg/L	Taste/odor, laxative effect for people not used to the water
	For more information, about Maximum Contentinent Levels (MCLs), health affects, and the threather the order of the contentinent and many			


For more information about Maximum Contaminant Levels (MCLs), health effects, and treatment options for these contaminants and more, see the NDDEQ's fact sheets (deq.nd.gov/wq/1_Groundwater) or visit the US EPA website (epa.gov/ground-water-and-drinking-water).

Dominant Water Type	Water Hardness
Calcium-Bicarbonate	Very Hard

Nitrate

Percentage of Wells Exceeding the Nitrate Maximum Contaminant Level (MCL)* (10 mg/L as N).

No Nitrate MCL Exceedances



Pesticides

Percentage of wells with detections of each pesticide detected in the aquifer.

No Pesticide Detections

State Pesticide Management Plan

Agricultural Groundwater Monitoring Program aquifers are monitored as a part of the State Pesticide Management Plan. A Prevention Action Level (PAL) threshold of 25% of the pesticide's Maximum Contaminant Level (MCL)* or Health Advisory Level (HAL) is used to identify whether action is needed to prevent further contamination.

Prevention Action Level Exceedances	None
MCL or HAL Exceedances	None

Number of Unique Wells with Pesticide Detections since 1996

0 of 8 Total Wells

2021 Pesticide Detections

No Pesticide Detections